


RG2833

Cat. No.:	HY-16425		
CAS No.:	1215493-56-3		
Molecular Formula:	C ₂₀ H ₂₅ N ₃ O ₂		
Molecular Weight:	339.43		
Target:	HDAC		
Pathway:	Cell Cycle/DNA Damage; Epigenetics		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 50 mg/mL (147.31 mM)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg
	Concentration				
	1 mM		2.9461 mL	14.7306 mL	29.4612 mL
	5 mM		0.5892 mL	2.9461 mL	5.8922 mL
	10 mM		0.2946 mL	1.4731 mL	2.9461 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (7.37 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (7.37 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (7.37 mM); Clear solution
- Add each solvent one by one: 5% DMSO >> 40% PEG300 >> 5% Tween-80 >> 50% saline
Solubility: ≥ 2.5 mg/mL (7.37 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

RG2833 is a brain-penetrant HDAC inhibitor with IC₅₀s of 60 nM and 50 nM for HDAC1 and HDAC3, respectively. The K_i values for HDAC1 and HDAC3 are 32 and 5 nM, respectively^[1].

IC₅₀ & Target

HDAC3	HDAC1	HDAC3	HDAC1
-------	-------	-------	-------

	50 nM (IC ₅₀)	60 nM (IC ₅₀)	5 nM (Ki)	32 nM (Ki)
In Vitro	<p>The K_i values of RG2833 for HDAC1 and HDAC3 are 32 nM and 5 nM, respectively. RG2833 is highly active in the whole tested concentration range from 1 to 10 μM. Continuous incubation with RG2833 slows the increase in frataxin protein, and when the compound is removed, frataxin protein levels rapidly increased in the cells from patient P13^[1]. RG2833 produces significant increases in brain aconitase enzyme activity, together with reduction of neuronal pathology of the dorsal root ganglia (DRG)^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>			
In Vivo	<p>RG2833 (150 mg/kg) is able to correct frataxin deficiency in the brain and heart of KIKI mice 24 hours after a single injection, but not when lower doses are used. When followed in time, the frataxin mRNA increase induced by RG2833 in the KIKI mouse can be first detected at 12 hours and reach a maximum at 24 hours in both brain and heart^[1]. RG2833 (100 mg/kg, s.c.) is well tolerated in chronic dosing of mice without toxicity. RG2833 improves motor coordination of YG8R FRDA mice. RG2833 increases frataxin protein expression in the brain of YG8R FRDA mice^[2]. RGFP109 (30 mg/kg, p.o. once daily for 6 days) has no acute effects on dyskinesia after single or 6 days once-daily treatment. One week following cessation of RGFP109, dyskinesia and duration of ON-time with disabling dyskinesia are reduced by 37% and 50%, respectively^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>			

PROTOCOL

Kinase Assay ^[2]

Aconitase activities are determined by homogenization of mouse brain tissues on ice at 10% w/v in CellLytic MT Mammalian Tissue Lysis/Extraction buffer, followed by centrifugation at 800×g for 10 min at 4°C. Tissue lysates (50 μL) are then added to 200 μL of substrate mix (50 mM Tris/HCl pH 7.4, 0.4 mM NADP, 5 mM Na citrate, 0.6 mM MgCl₂, 0.1% (v/v) Triton X-100 and 1U isocitrate dehydrogenase) and the reactions are incubated at 37°C for 15 min, followed by spectrophotometric absorbance measurements every minute for 15 min at 340 nm 37°C to determine the reaction slope. Aconitase activities of mouse brain tissues are then normalized to citrate synthase activities, which are determined using a citrate synthase assay kit.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[2]

Mice are housed in conventional open cages with Litaspen Premium 8/20 bedding, paper wool nesting and standard fun tunnel environmental enrichment, with 13 h light, 11 h dark, 20-23°C and 45-60% humidity. The mice are given a diet of SDS RM3 Expanded food pellets and standard drinking water. Mice are given subcutaneous injections of 150 mg/kg RG2833 three times per week for 4.5 months, or 50 mg/kg 136 or 100 mg/kg RG2833 five times per week for 5 months, followed by culling for tissue collection 24 h after the final injection.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cancer Res. 2021 Dec 15;81(24):6131-6141.
- Acta Pharmacol Sin. 2021 Apr 13.
- J Mol Med (Berl). 2019 Aug;97(8):1183-1193.
- ACS Med Chem Lett. 2015 Jun 22;6(8):948-52.
- PLoS One. 2016 Apr 14;11(4):e0153574.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Rai M, et al. Two new pimelic diphenylamide HDAC inhibitors induce sustained frataxin upregulation in cells from Friedreich's ataxia patients and in a mouse model. PLoS One. 2010, 5(1), e8825.

[2]. Sandi C, et al. Prolonged treatment with pimelic o-aminobenzamide HDAC inhibitors ameliorates the disease phenotype of a Friedreich ataxia mouse model. Neurobiol Dis. 2011, 42(3), 496-505.

[3]. Johnston TH, et al. RGFP109, a histone deacetylase inhibitor attenuates L-DOPA-induced dyskinesia in the MPTP-lesioned marmoset: a proof-of-concept study. Parkinsonism Relat Disord. 2013, 19(2), 260-264.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA