


trans-Zeatin

Cat. No.:	HY-19700		
CAS No.:	1637-39-4		
Molecular Formula:	C ₁₀ H ₁₃ N ₅ O		
Molecular Weight:	219.24		
Target:	MEK; ERK; Endogenous Metabolite		
Pathway:	MAPK/ERK Pathway; Stem Cell/Wnt; Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (114.03 mM; ultrasonic and warming and heat to 60°C)					
	Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg
		Concentration				
		1 mM		4.5612 mL	22.8061 mL	45.6121 mL
		5 mM		0.9122 mL	4.5612 mL	9.1224 mL
10 mM		0.4561 mL	2.2806 mL	4.5612 mL		
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1.25 mg/mL (5.70 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1.25 mg/mL (5.70 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 1.25 mg/mL (5.70 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	trans-Zeatin is a plant cytokinin, which plays an important role in cell growth, differentiation, and division; trans-Zeatin also inhibits UV-induced MEK/ERK activation.	
IC₅₀ & Target	MEK	ERK
In Vitro	trans-Zeatin is a plant cytokinin, which plays an important role in cell growth, differentiation, and division ^[1] . trans-Zeatin (20, 40 or 80 μM) inhibits UV-induced MEK/ERK activation, upregulates AQP3 in a time- and dose-dependent manner, and attenuates UV induced loss of AQP3 in keratinocytes (HaCaT cells). UV-induced AQP3 downregulation is blocked by MEK/ERK	

inhibitors. Trans-Zeatin (80 μ M) attenuates UV-induced downregulation of wound healing and water permeability in HaCaT cells^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nat Plants. 2024 Jan;10(1):180-191.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Li Q, et al. Endogenous trans-zeatin content in plants with different metal-accumulating ability: a field survey. Environ Sci Pollut Res Int. 2016 Dec;23(23):23422-23435. Epub 2016 Sep 9.

[2]. Ji C, et al. Trans-Zeatin attenuates ultraviolet induced down-regulation of aquaporin-3 in cultured human skin keratinocytes. Int J Mol Med. 2010 Aug;26(2):257-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA