


## AZ7550 Mesylate

Cat. No.:	HY-B0794B
CAS No.:	2319837-99-3
Molecular Formula:	C <sub>30</sub> H <sub>43</sub> N <sub>7</sub> O <sub>11</sub> S <sub>3</sub>
Molecular Weight:	773.9
Target:	IGF-1R; Drug Metabolite; EGFR
Pathway:	Protein Tyrosine Kinase/RTK; Metabolic Enzyme/Protease; JAK/STAT Signaling
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (32.30 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg	
				1 mM	1.2922 mL	6.4608 mL	12.9216 mL
				5 mM	0.2584 mL	1.2922 mL	2.5843 mL
10 mM				0.1292 mL	0.6461 mL	1.2922 mL	
Please refer to the solubility information to select the appropriate solvent.							
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (3.23 mM); Clear solution						
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (3.23 mM); Clear solution						

### BIOLOGICAL ACTIVITY

Description	AZ7550 Mesylate is an active metabolite of AZD9291 and inhibits the activity of IGF1R with an IC <sub>50</sub> of 1.6 μM.
IC <sub>50</sub> & Target	IC <sub>50</sub> : 1.6 μM (IGF1R), 88 nM (MLK1), 156 nM (ACK1), 195 nM (ErbB4), 228 nM (MNK2), 302 nM (FLT3), 420 nM (ALK), 449 nM (FES), 840 nM (IRR), 843 nM (BRK), 977 nM (BLK), 995 nM (FAK), 1256 nM (Ins R), 1317 nM (TEC), 1784 nM (FLT4), 2288 nM (PYK2), 2443 nM (Txk), 5104 nM (BTK) <sup>[1]</sup>
In Vitro	AZ7550 (Compound 28) appears to offer a broadly similar potency and selectivity profile to the parent compound AZD9291. AZ7550 inhibits double mutant (DM) cell line H1975, activating mutant (AM) cell line PC9, and wild type (WT) cell line LoVo with IC <sub>50</sub> s of 45, 26, and 786 nM, respectively. AZ7550 inhibits DM antiproliferative cell line H1975, AM antiproliferative cell line PC9, and WT antiproliferative cell line Calu3 with GI <sub>50</sub> s of 19, 15, and 537 nM, respectively <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## PROTOCOL

### Kinase Assay <sup>[1]</sup>

Biochemical enzyme profiling of AZD9291 and active metabolites across the kinome panel (single profiling experiment representative of two independent studies). % inhibition for kinases in the ~280 kinase panel that shows greater than 60% inhibition after 1  $\mu$ M treatment with AZD9291, AZ5104 or AZ7550, and follow-up IC<sub>50</sub>s where tested, are shown. Kinases with a conserved cysteine in the analogous position within their catalytic domain as Cys797 in EGFR are also shown, highlighted in bold<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Biopharm Drug Dispos. 2023 Jan 17.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Finlay MR, et al. Discovery of a potent and selective EGFR inhibitor (AZD9291) of both sensitizing and T790M resistance mutations that spares the wild type form of the receptor. J Med Chem. 2014 Oct 23;57(20):8249-67.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA