


Acetylcholine-d₉ chloride

Cat. No.:	HY-B0282S1
CAS No.:	344298-95-9
Molecular Formula:	C ₇ H ₇ D ₉ ClNO ₂
Molecular Weight:	190.72
Target:	nAChR; Calcium Channel; Endogenous Metabolite
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (524.33 mM; Need ultrasonic)
DMSO : 62.5 mg/mL (327.71 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	5.2433 mL	26.2164 mL	52.4329 mL
	5 mM	1.0487 mL	5.2433 mL	10.4866 mL
	10 mM	0.5243 mL	2.6216 mL	5.2433 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Acetylcholine-d₉ (chloride) is the deuterium labeled Acetylcholine chloride. Acetylcholine chloride (ACh chloride), a neurotransmitter, is a potent cholinergic agonist. Acetylcholine chloride is a modulator of the activity of dopaminergic (DAergic) neurons through the stimulation of nicotinic acetylcholine receptors (nAChRs)[1][2]. Acetylcholine chloride inhibits p53 mutant peptide aggregation in vitro[5].

In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019;53(2):211-216.
- [2]. Prashant Tiwari, et al. Basic and modern concepts on cholinergic receptor: A review. *Asian Pac J Trop Dis*. 2013 Oct;3(5): 413-420.

-
- [3]. A Young, et al. Diarrhoea of famine and malnutrition--investigations using a rat model. 2--Ileal hypersecretion induced by starvation. Gut. 1990 Feb;31(2):162-9.
- [4]. Xia Lei, et al. Effects of acetylcholine chloride on intracellular calcium concentration of cultured sweat gland epithelial cells. Arch Dermatol Res. 2008 Aug;300(7):335-41.
- [5]. P F Zabrodskii, et al. Effect of acetylcholine on mortality of mice from sepsis and proinflammatory cytokine production. Bull Exp Biol Med. 2011 Jan;150(3):340-2.
- [6]. Zhaolin Chen, et al. Inhibition of p53 Mutant Peptide Aggregation In Vitro by Cationic Osmolyte Acetylcholine Chloride. Protein Pept Lett. 2017;24(4):353-357.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA