


Amoxicillin arginine

Cat. No.:	HY-B0467C
CAS No.:	59261-05-1
Molecular Formula:	C ₁₆ H ₁₉ N ₃ O ₅ S.C ₆ H ₁₄ N ₄ O ₂
Target:	Antibiotic; Bacterial
Pathway:	Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Amoxicillin (Amoxycillin) arginine is an antibiotic with good oral absorption and broad spectrum antimicrobial activity. Amoxicillin arginine inhibits the biosynthesis of polypeptides in the cell wall, thereby inhibiting cell growth ^{[1][2][3]} .
In Vitro	Amoxicillin (Amoxycillin) arginine (1-100 μM; 24 hours; <i>L. acidophilus</i>) decreases living cells and increases degree of cell wall rupture in a dose-dependent manner ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Amoxicillin (Amoxycillin) arginine (7 mg/kg; i.h.; female ICR/Swiss mice) inhibits strain numbers and improves the survival rate of rats in 1 mg/L or less ^[2] . Amoxicillin (Amoxycillin) arginine (1.6-9.5 mg/kg; p.o.; daily, for 7 or 14 days; swiss albino mice) has against infection with chlamydia trachomatis in mice ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
	Animal Model: Female ICR/Swiss mice ^[2]
	Dosage: 7 mg/kg
	Administration: Subcutaneous injection; every 8 h, for 24 hours
	Result: Inhibited bacterial numbers in a dose-dependent manner.
	Animal Model: Female ICR/Swiss mice ^[2]
	Dosage: 7 mg/kg
	Administration: Subcutaneous injection; every 8 h, for 4 days
	Result: Survived all animals that were infected with organisms for which MICs were 1 mg/L or less, and with the two strains for which MICs were 2 mg/L, 20 to 40% mortality.
Animal Model: Swiss albino mice ^[3]	

Dosage:	1.6 and 9.5 mg/kg
Administration:	Oral administration; daily, for 7 or 14 days
Result:	Improved the activity of Chlamydia trachomatis infection in mice.

CUSTOMER VALIDATION

- Nat Commun. 2022 Mar 2;13(1):1116.
- Chemosphere. 2023 Oct 3;344:140353.
- Chemosphere. 2019 Jun;225:378-387.
- Environ Sci Pollut Res Int. 2017 Feb;24(6):5918-5932.
- Antimicrob Agents Chemother. 2021 Feb 17;65(3):e01921-20.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Guo Y, et, al. Metabolic response of Lactobacillus acidophilus exposed to amoxicillin. J Antibiot (Tokyo). 2022 May;75(5):268-281.
- [2]. Andes D, et, al. In vivo activities of amoxicillin and amoxicillin-clavulanate against Streptococcus pneumoniae: application to breakpoint determinations. Antimicrob Agents Chemother. 1998 Sep;42(9):2375-9.
- [3]. Kramer MJ, et, al. Activity of oral amoxicillin, ampicillin, and oxytetracycline against infection with chlamydia trachomatis in mice. J Infect Dis. 1979 Jun;139(6):717-9.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA