


Bestatin trifluoroacetate

Cat. No.:	HY-B0134B
CAS No.:	223763-80-2
Molecular Formula:	C ₁₈ H ₂₅ F ₃ N ₂ O ₆
Molecular Weight:	422.4
Target:	Aminopeptidase; Bacterial; Antibiotic
Pathway:	Metabolic Enzyme/Protease; Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Bestatin trifluoroacetate is an inhibitor of CD13 (Aminopeptidase N)/APN and leukotriene A4 hydrolase, used for cancer research.
IC₅₀ & Target	CD13
In Vitro	<p>Bestatin enhances ATRA-induced differentiation and inhibits ATRA-driven phosphorylation of p38 MAPK in ATRA-sensitive APL NB4 cells. Bestatin can not reverse the differentiation block in ATRA-resistant APL MR2 cells. CD13 ligation with anti-CD13 antibody WM-15 results in phosphorylation of p38 MAPK, reduces the inhibition of Bestatin on the phosphorylation of p38 MAPK, and completely abolishes the enhancement of Bestatin on ATRA-inducing differentiation in NB4 cells^[2]. Bestatin (600 μM)-treated cells progress slower through the cell cycle due to decreased rate of cell growth and the frequency of cell division. Bestatin inhibits the frequency of mitosis and the inherent multinuclearity in <i>D. discoideum</i>, and is not cytotoxic to <i>D. discoideum</i> cells at 0-600 μM. Bestatin inhibits aminopeptidase activity in lysates of PsaA-GFP- and GFP-expressing cells by 69.39% ± 10.5% and 39.93% ± 18.7% of control, respectively^[4].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
In Vivo	<p>Bestatin (20 μM) significantly reduces CD13 expression in diabetic mice and results a significant inhibition of MMP-9 specific gelatinolytic band densities compared to diabetic vehicle-treated mice. Bestatin treatment significantly inhibits the expression of VEGF and heparanase in diabetic mice. Intravitreal bestatin treatment significantly downregulates the expression of both HIF-1α and VEGF in diabetic mice retinas. Furthermore, the upregulated expression of heparanase in diabetic mice retinas is significantly inhibited by intravitreal bestatin treatment^[1]. Bestatin (10, 1, and 0.1mg/kg, i.p.) treatment before the antigen-potentiated humoral response to SRBC results in an increased number of splenocytes producing hemolytic anti-SRBC antibodies (PFC) and the 2-ME-resistant serum hemagglutinin titer (at a dose of 0.1 mg/kg). Bestatin (1 and 0.1 mg/kg) administered to mice five times on alternate days after cyclophosphamide injection does not change the suppressive effect of the drug regarding the number of PFC, and even causes the further decrease of the total anti-SRBC hemagglutinins at dose of 1 mg/kg on day 7 after antigen stimulation^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

PROTOCOL

Kinase Assay ^[4]

Cells are harvested, washed, and lysed in NP-40 lysis buffer (50 mM Tris-HCl [pH 7.5], 150 mM NaCl, 0.5% NP-40). Total cell protein is quantified using the Bradford assay and 1-mg/mL protein aliquots are made. Ten microliters of total cell protein is

	<p>mixed with 290 μL of substrate solution (0.1 mg/mL dithiothreitol [DTT], 0.1 mg/mL albumin, and 1 mM alanine-β-naphthylamide). Fluorometric measurements (340 nm excitation, 400 nm emission) are made after 15 and 30 min. The slope of the line between the 15- and 30-min measurements is used to represent aminopeptidase activity. Total cell protein is preincubated with bestatin, amastatin, puromycin, EDTA, and/or ZnCl₂ for 20 min before the fluorometric aminopeptidase assay.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Cell Assay ^[4]	<p>Growing cells (1\times10⁶ to 2\times10⁶ cells/mL) are diluted to 1.0\times10³ cells/mL and transferred (3 mL) into a well in a 12-well multiwell plate (2.5-cm diameter/well). Cells are treated with 0, 10, 50, 100, 300, or 600 μM Bestatin and allowed to grow at 21°C shaking at 180 rpm for 48 h. A hemocytometer is used to measure cell density after 0, 24, and 48 h.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Animal Administration ^[3]	<p>Bestatin is dissolved in PBS. The agent (doses of 10, 1, and 0.1 mg/kg) is injected i.p. to non-cyclophosphamide-treated mice, 5 or 10 times at 24-h intervals before SRBC immunization. The mice are immunized 24 h after the last dose of bestatin. Pharmacological immunosuppression is induced by a single intraperitoneal injection of cyclophosphamide administered at a dose of 350 mg/kg, 12 days before SRBC immunization. Bestatin at the doses of 1 and 0.1 mg/kg is injected to cyclophosphamide-immunosuppressed mice i.p. five times at 48-h intervals or 10 times at 24-h intervals before SRBC immunization. The first dose of bestatin is administered 24 h after cyclophosphamide, while the last dose of the drug is injected 24h before SRBC immunization.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

CUSTOMER VALIDATION

- J Med Chem. 2017 Mar 9;60(5):1817-1828.
- Int J Oncol. 2019 Jul;55(1):331-339.
- J Pharmacol Exp Ther. 2022 Aug;382(2):188-198.
- Dis Markers. 11 Jan 2022.
- Hemasphere. 2021 Jun 12;5(7):e602.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Hossain A, et al. Protective effects of bestatin in the retina of streptozotocin-induced diabetic mice. *Exp Eye Res.* 2016 Aug;149:100-6
- [2]. Qian X, et al. Inhibition of p38 MAPK Phosphorylation Is Critical for Bestatin to Enhance ATRA-Induced Cell Differentiation in Acute Promyelocytic Leukemia NB4 Cells. *Am J Ther.* 2016 May-Jun;23(3):e680-9.
- [3]. Lis M, et al. The effects of bestatin on humoral response to sheep erythrocytes in non-treated and cyclophosphamide-immunocompromised mice. *Immunopharmacol Immunotoxicol.* 2013 Feb;35(1):133-8
- [4]. Poloz Y, et al. Bestatin inhibits cell growth, cell division, and spore cell differentiation in *Dictyostelium discoideum*. *Eukaryot Cell.* 2012 Apr;11(4):545-57

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA