


## CY5-SE

<b>Cat. No.:</b>	HY-D0819
<b>CAS No.:</b>	146368-14-1
<b>Molecular Formula:</b>	C <sub>37</sub> H <sub>43</sub> N <sub>3</sub> O <sub>10</sub> S <sub>2</sub>
<b>Molecular Weight:</b>	753.88
<b>Target:</b>	Fluorescent Dye
<b>Pathway:</b>	Others
<b>Storage:</b>	-20°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



## SOLVENT & SOLUBILITY

### In Vitro

DMSO : 25 mg/mL (33.16 mM; Need ultrasonic)  
 H<sub>2</sub>O : ≥ 5.88 mg/mL (7.80 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	1.3265 mL	6.6324 mL	13.2647 mL
	5 mM	0.2653 mL	1.3265 mL	2.6529 mL
	10 mM	0.1326 mL	0.6632 mL	1.3265 mL

Please refer to the solubility information to select the appropriate solvent.

### In Vivo

- Add each solvent one by one: Saline  
Solubility: 25 mg/mL (33.16 mM); Clear solution; Need ultrasonic and warming and heat to 60°C
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.5 mg/mL (3.32 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (3.32 mM); Clear solution

## BIOLOGICAL ACTIVITY

### Description

Cy5-SE (Cy5 NHS Ester) is a reactive dye for the labeling of amino-groups in peptides, proteins, and oligonucleotides. This dye requires small amount of organic co-solvent (such as DMF or DMSO) to be used in labeling reaction. This reagent is ideal for very cost-efficient labeling of soluble proteins, as well as all kinds of peptides and oligonucleotides. This reagent also works well in organic solvents for small molecule labeling. Excitation (nm): 649, Emission (nm): 670.

### In Vitro

Guidelines (Following is our recommended protocol. This protocol only provides a guideline, and should be modified)

according to your specific needs).

Conjugation of SELP Analogues with Cy5-SE.

1. 1 mL of each 100 µg/mL SELP solution in 100 mM sodium bicarbonate buffer (pH 8.3) is mixed with 10 µL of 1.2 mg/mL Cy5 mono NHS-ester in 10% DMSO and incubates for 2 hours on ice.

2. To quench the reaction, 50 µL of 1 M Tris-HCl (pH 8.0) is added to the reaction solution.

3. Reaction mixtures are loaded onto 1.5 mL Sephadex G-25 columns, and Cy5-conjugated SELPs are eluted by centrifugation for 3 minutes at 1050g<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Adv Funct Mater. 2023 Sep 15.
- Adv Sci (Weinh). 2023 Jun 25;e2301592.
- J Nanobiotechnology. 2018 Mar 16;16(1):23.
- Nano Res. 29 June 2021.
- ACS Appl Mater Interfaces. 2019 Jan 16;11(2):1766-1781.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Jeon HY, et al. Array-Based High-Throughput Analysis of Silk-Elastinlike Protein Polymer Degradation and C-Peptide Release by Proteases. Anal Chem. 2016;88(10):5398-5405.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA