Proteins

Canthaxanthin

Cat. No.: HY-B1960 CAS No.: 514-78-3 Molecular Formula: $C_{40}H_{52}O_2$ Molecular Weight: 564.84

Target: Reactive Oxygen Species; Endogenous Metabolite

Pathway: Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κΒ

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

Product Data Sheet

BIOLOGICAL ACTIVITY

| Description | Canthaxanthin is a red-orange carotenoid with various biological activities, such as antioxidant, antitumor properties. |
|---------------------------|--|
| IC ₅₀ & Target | Human Endogenous Metabolite |
| In Vitro | Canthaxanthin enrichment of LDL has the potential to protect cholesterol from oxidation. In addition to its free radical scavenging and antioxidant properties (e.g., the induction of catalase and superoxide dismutase), canthaxanthin shows immunomodulatory activity (e.g. enhancing the proliferation and function of immune competent cells) and plays important role in gap junction communication (e.g. induction of the transmembrane protein connexin) ^[1] . At concentrations of 0.1 to 1 x 1000 μ M, canthaxanthin significantly reduces the overall number of tumor cells. The greatest inhibition is observed at a canthaxanthin concentration of 1000 after 72 h and 96 h of incubation ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. |
| In Vivo | Canthaxanthin alters the protective ability of tissues against oxidative stress. Canthaxanthin treatment for 15 d at the dose of 14 μ g/kg body weight results in hepatic incorporation of the carotenoid, which is maximum in liver and reaches 0.52 \pm 0.05 nmol/g liver. Glutathione peroxidase activity is 35% lower and catalase (59%) and manganese superoxide dismutase (28%) activities are higher in canthaxanthin-treated mice than in controls ^[3] . Canthaxanthin? inhibit the growth of mammary tumors in mice and the anti-tumor activity is also influenced by the supplemental dose ^[4] . Diet supplementation with canthaxanthin for 3 weeks prior to the carcinogen results in a 65% reduction in the number of mammary cancers by a mechanism not involving pro-vitamin A activity ^[5] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. |

PROTOCOL

Animal Administration [3] Mice: Female 6-wk-old Balb/c mice are randomly divided into two groups (n = 10/group). The control group receives olive oil alone (vehicle) and the canthaxanthin-treated group receives canthaxanthin at a dose of 14 µg/kg body weight per day. At the end of the treatment, mice are killed by cervical dislocation and liver is excised, frozen in liquid nitrogen and stored at

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- FASEB J. 2020 Oct;34(10):13430-13444.
- Biochem Biophys Res Commun. 2018 Sep 3;503(1):297-303.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Esatbeyoglu T, et al. Canthaxanthin: From molecule to function. Mol Nutr Food Res. 2017 Jun;61(6).
- [2]. Huang DS, et al. Inhibitory effects of canthaxanthin on in vitro growth of murine tumor cells. Cancer Lett. 1992 Aug 31;65(3):209-13.
- [3]. Palozza P, et al. Canthaxanthin supplementation alters antioxidant enzymes and iron concentration in liver of Balb/c mice. J Nutr. 2000 May;130(5):1303-8.
- [4]. Chew BP, et al. A comparison of the anticancer activities of dietary beta-carotene, canthaxanthin and astaxanthin in mice in vivo. Anticancer Res. 1999 May-Jun;19(3A):1849-53.
- [5]. Grubbs CJ, et al. Effect of canthaxanthin on chemically induced mammary carcinogenesis. Oncology. 1991;48(3):239-45.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA