


Cefmenoxime

Cat. No.:	HY-B0875A
CAS No.:	65085-01-0
Molecular Formula:	C ₁₆ H ₁₇ N ₉ O ₅ S ₃
Molecular Weight:	511.56
Target:	Antibiotic; Bacterial
Pathway:	Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Cefmenoxime (SCE-1365) is a new semisynthetic cephalosporin antibiotic. Cefmenoxime has antibacterial activity against a wide variety of gram-positive and gram-negative bacteria ^{[1][2]} .								
IC₅₀ & Target	β-lactam								
In Vitro	<p>Cefmenoxime (SCE-1365) inhibits at least 90% of strains tested (MIC₉₀) ranged from 0.06 to 8 μg/mL for the Enterobacteriaceae^[1].</p> <p>Cefmenoxime (SCE-1365) inhibits MIC₉₀ values for gram-positive cocci are 0.015 and ≤0.008 μg/mL for Streptococcus pneumoniae and Streptococcus pyogenes, respectively, and 2 μg/mL for S. aureus^[1].</p> <p>Cefmenoxime (SCE-1365) against Haemophilus influenzae, Neisseria gonorrhoeae and Neisseria meningitidis with MIC₉₀ values ranging from ≤0.008 to 0.25 μg/mL^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>								
In Vivo	<p>Cefmenoxime (SCE-1365) (40 mg/kg; i.h.; daily, for 7 d; male Jcl:ICR mice) improves the survival rate of mice infected with lung bacteria^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;">Animal Model:</td> <td>Male Jcl:ICR mice^[2]</td> </tr> <tr> <td>Dosage:</td> <td>40 mg/kg</td> </tr> <tr> <td>Administration:</td> <td>Subcutaneous injection; daily, for 7 days</td> </tr> <tr> <td>Result:</td> <td>Inhibited mortality rate of animals was 60% at a dose of 40 mg/kg.</td> </tr> </table>	Animal Model:	Male Jcl:ICR mice ^[2]	Dosage:	40 mg/kg	Administration:	Subcutaneous injection; daily, for 7 days	Result:	Inhibited mortality rate of animals was 60% at a dose of 40 mg/kg.
Animal Model:	Male Jcl:ICR mice ^[2]								
Dosage:	40 mg/kg								
Administration:	Subcutaneous injection; daily, for 7 days								
Result:	Inhibited mortality rate of animals was 60% at a dose of 40 mg/kg.								

REFERENCES

[1]. Stamm JM, et, al. Antimicrobial activity of cefmenoxime (SCE-1365). Antimicrob Agents Chemother. 1981 Mar;19(3):454-60.

[2]. Tatara O, et, al. Synergistic effects of romurtide and cefmenoxime against experimental Klebsiella pneumonia in mice. Antimicrob Agents Chemother. 1992 Jan;36(1):167-71.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA