Cefmetazole sodium

MedChemExpress

®

Cat. No.:	HY-B1257	
CAS No.:	56796-39-5	
Molecular Formula:	$C_{15}H_{16}N_7NaO_5S_3$	N ^N N O ONa
Molecular Weight:	493.52	N= S S
Target:	Bacterial; Antibiotic	S S N
Pathway:	Anti-infection	H O H
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)	

SOLVENT & SOLUBILITY

S		Solvent Mass Concentration	1 mg	5 mg	10 mg	
	Preparing Stock Solutions	1 mM	2.0263 mL	10.1313 mL	20.2626 mL	
		5 mM	0.4053 mL	2.0263 mL	4.0525 mL	
		10 mM	0.2026 mL	1.0131 mL	2.0263 mL	
	Please refer to the so	lubility information to select the app	propriate solvent.			
In Vivo		1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (4.21 mM); Clear solution				
		2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (4.21 mM); Clear solution				

BIOLOGICAL ACTIVITY				
Description	Cefmetazole sodium (Sodium cefmetazole) is a semisynthetic cephamycin antibiotic with broad-spectrum antibacterial activity, covering gram-positive, gram-negative, and anaerobic bacteria. Cefmetazole sodium binds to penicillin binding proteins (PBPs), resulting in interfering bacterial cell wall biosynthesis. Cefmetazole sodium is used for the research of gynecologic, intraabdominal, urinary tract, respiratory tract and skin and soft tissue infections ^{[1][2][3]} .			
IC ₅₀ & Target	β-lactam			

Product Data Sheet

In Vitro	Cefmetazole sodium (Sodium cefmetazole) has antibiotic affinity to S. aureus with a MIC value of 1.0 mg/L and has affinity for PBP1, PBP2 and PBP3 with IC ₅₀ values of ≤0.3, 0.109 and 0.494 mg/L ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.		
In Vivo	Cefmetazole sodium (Sodium cefmetazole) (100 mg/kg; i.h.; twice a day, for 7 days; male ICR mice) alters gut bacterial flora [3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only. Animal Model: Male ICR mice ^[3]		
	Dosage:	100 mg/kg	
	Administration:	Subcutaneous injection; twice a day, for 7 days	
	Result:	Reduced Peyer's patch (PP) lymphocyte cell numbers while decreased bacterial numbers in the small intestine.	

REFERENCES

[1]. Truesdell SE, et, al. Interaction of cephalosporins with penicillin-binding proteins of methicillin-resistant Staphylococcus aureus. J Antimicrob Chemother. 1989 Apr;23 Suppl D:13-9.

[2]. Yaguchi Y, et, al. Influences of long-term antibiotic administration on Peyer's patch lymphocytes and mucosal immunoglobulin A levels in a mouse model. JPEN J Parenter Enteral Nutr. 2006 Sep-Oct;30(5):395-8; discussion 399.

[3]. Schentag JJ, et, al. Cefmetazole sodium: pharmacology, pharmacokinetics, and clinical trials. Pharmacotherapy. 1991;11(1):2-19.

Caution: Product has not been fully validated for medical applications. For research use only.