


Chloroxine

Cat. No.:	HY-B0295		
CAS No.:	773-76-2		
Molecular Formula:	C ₉ H ₅ Cl ₂ NO		
Molecular Weight:	214		
Target:	Bacterial; Antibiotic; Parasite		
Pathway:	Anti-infection		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : 33.33 mg/mL (155.75 mM; Need ultrasonic)
 H₂O : < 0.1 mg/mL (insoluble)

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		4.6729 mL	23.3645 mL	46.7290 mL
	5 mM		0.9346 mL	4.6729 mL	9.3458 mL
	10 mM		0.4673 mL	2.3364 mL	4.6729 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: 2.5 mg/mL (11.68 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: 2.5 mg/mL (11.68 mM); Suspended solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

Chloroxine is one of the important 8-hydroxyquinoline derivative. Chloroxine has effective antibacterial, antifungal, antiprotozoal and antiamebic activities, especially used in treating the intestinal amebiasis. Chloroxine is also used in the treatment of dandruff and seborrheic dermatitis of the scalp^{[1][2]}.

IC₅₀ & Target

Amebae

In Vitro

By using the Jouyban-Acree model, the Chloroxine solubility is well correlated obtaining RAD values lower than 3.64% and RMSD values lower than 8.82×10^{-6} . Chloroxine is preferentially solvated by water for the studied mixtures in water-rich compositions; while within intermediate and co-solvent-rich compositions, Chloroxine is preferentially solvated by DMSO

(DMF, NMP or 1,4-dioxane) in DMSO (DMF, NMP or 1,4-dioxane) + water mixtures. It is shown that the change in solvent-solvent interaction energy accounted by cavity term governed the solubility variation of Chloroxine in all aqueous mixtures [1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Clin Transl Oncol. 2023 Oct 18.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Renjie X, et al. Solubility modelling, solvent effect and preferential solvation of carbendazim in aqueous co-solvent mixtures of, N,N -dimethylformamide, methanol, ethanol and, n -propanol. J. Chem. Thermodynamics 138 (2019) 288–296.

[2]. Chloroxine (Topical). Dec 27, 2018

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA