

Crocetin meglumine

Cat. No.: HY-42937

Molecular Formula: $C_{34}H_{58}N_2O_{14}$


Molecular Weight: 718.83

Target: iGluR

Pathway: Membrane Transporter/Ion Channel; Neuronal Signaling

Storage: 4°C, sealed storage, away from moisture

* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 8.33 mg/mL (11.59 mM); ultrasonic and warming and heat to 66°C

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	1.3911 mL	6.9557 mL	13.9115 mL
	5 mM	0.2782 mL	1.3911 mL	2.7823 mL
	10 mM	0.1391 mL	0.6956 mL	1.3911 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: 0.83 mg/mL (1.15 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE- β -CD in saline)
Solubility: \geq 0.83 mg/mL (1.15 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Crocetin (Transcrocetin) meglumine, extracted from saffron (*Crocus sativus* L.), acts as an NMDA receptor antagonist with high affinity.

IC₅₀ & Target

NMDA Receptor

In Vitro

Crocetin (Transcrocetin) meglumine, a saffron metabolite originating from the crocin apocarotenoids, has been shown to exert strong NMDA receptor affinity and is thought to be responsible for the CNS activity of saffron. To ensure unchanged viability of Caco-2 cells throughout the transport experiments, cellular mitochondrial dehydrogenase activity of Caco-2 cells is measured by MTT assay after a 24 h incubation period with the test compounds: Hydroalcoholic saffron extract saffron extract (SE, 0.5-1 mg/mL) and crocin-1 (250-1000 μ M) reveal no negative significant changes in cellular viability. Crocetin meglumine at 10 μ M level does not change viability while higher concentrations (40-160 μ M) reduces significantly cellular

viability^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

Cytotoxicity of test compounds is determined by MTT assay using Caco-2 cells in 96 well plates at a density of 20,000 cells per well in 200 µl FBS-free medium, grown for 96 h and followed by 24 h contact time with the test compounds (100 µL of serum-free media containing SE 0.5, 1, and 2 mg/mL; trans-crocin-1 250, 500, and 1000 µM; Transcrocetin 10, 40, 80, and 160 µM) and incubation at 37°C/5% CO₂. The incubation solutions are aspirated, each well is washed twice with 150 µL of PBS and 50 µL of MTT solution are added (2.5 mg/mL in PBS). Supernatants are discarded and the formed formazan is dissolved in 50 µL of DMSO. The absorption of the resulting solution is determined at λ=492 nm against reference wavelength λ=690 nm^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Lautenschläger M, et al. Intestinal formation of trans-Crocetin from saffron extract (*Crocus sativus* L.) and in vitro permeation through intestinal and blood brain barrier. *Phytomedicine*. 2015 Jan 15;22(1):36-44.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA