Screening Libraries

Product Data Sheet

Ginkgolide A

Cat. No.: HY-B0355 CAS No.: 15291-75-5 Molecular Formula: $C_{20}H_{24}O_9$ Molecular Weight: 408.4

Target: GABA Receptor; Endogenous Metabolite

Pathway: Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease

Storage: Powder -20°C 3 years

4°C 2 years

In solvent -80°C 2 years

> -20°C 1 year

| \ HQ | - (T | 0 |
|-------------|------|-----|
| 0 | 0 | OH. |
| 0′ 1 | H — | |

SOLVENT & SOLUBILITY

In Vitro

DMSO: 100 mg/mL (244.86 mM; Need ultrasonic)

| Preparing Stock Solutions | Solvent Mass Concentration | 1 mg | 5 mg | 10 mg |
|------------------------------|-------------------------------|-----------|------------|------------|
| | 1 mM | 2.4486 mL | 12.2429 mL | 24.4858 mL |
| | 5 mM | 0.4897 mL | 2.4486 mL | 4.8972 mL |
| | 10 mM | 0.2449 mL | 1.2243 mL | 2.4486 mL |

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.12 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.12 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.12 mM); Clear solution

BIOLOGICAL ACTIVITY

| Description | Ginkgolide A (BN-52020) is an extract from in Ginkgo biloba and a g-aminobutyric acid (GABA) antagonist. |
|---------------------------|--|
| IC ₅₀ & Target | Human Endogenous Metabolite |
| In Vitro | Ginkgolide A (BN-52020) is a highly active PAF antagonist cage molecule that is isolated from the leaves of the Ginkgo biloba tree. Shows potential in a wide variety of inflammatory and immunological disorders. Ginkgolide A (BN-52020) failed to affect apoptotic damage neither in serum-deprived nor in staurosporine-treated neurons ^[2] . |

| | MCE has not independently confirmed the accuracy of these methods. They are for reference only. |
|---------|---|
| In Vivo | Ginkgolide A significantly shortened the sleeping time in anesthetized mice ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. |

CUSTOMER VALIDATION

- Life Sci. 2023 Jan 31;317:121439.
- Contrast Media Mol Imaging. 2022 Sep 10;2022:6384334.

See more customer validations on $\underline{www.MedChemExpress.com}$

REFERENCES

[1]. Wada, K., et al., Isolation of bilobalide and ginkgolide A from Ginkgo biloba L. shorten the sleeping time induced in mice by anesthetics. Biol Pharm Bull, 1993. 16(2): p. 210-2.

[2]. Ahlemeyer, B. and J. Krieglstein, Pharmacological studies supporting the therapeutic use of Ginkgo biloba extract for Alzheimer's disease. Pharmacopsychiatry, 2003. 36(S 1): p. 8-14.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA