


HKOH-1

Cat. No.:	HY-D1151
CAS No.:	2031170-96-2
Molecular Formula:	C ₂₆ H ₁₂ Cl ₂ I ₂ O ₆
Molecular Weight:	745.08
Target:	Reactive Oxygen Species; Fluorescent Dye
Pathway:	Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB; Others
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 180 mg/mL (241.58 mM; Need ultrasonic)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	1.3421 mL	6.7107 mL	13.4214 mL
5 mM	0.2684 mL	1.3421 mL	2.6843 mL
10 mM	0.1342 mL	0.6711 mL	1.3421 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

HKOH-1 is a highly sensitive green fluorescent probe for the specific detection of ·OH in living cells with a maximum excitation wavelength and emission wavelength of 500 nm and 520 nm, respectively^[1].

In Vitro

1. Preparation of HKOH-1 working solution
 - 1.1 Preparation of the stock solution
Dissolve 1 mg HKOH-1 in 135 μL DMSO to obtain 10 mM of stock solution.
Note: It is recommended to store the stock solution at -20℃ -80℃ away from light and avoid repetitive freeze-thaw cycles.
 - 1.2 Preparation of HKOH-1 working solution
Dilute the stock solution in serum-free cell culture medium or PBS to obtain 1-10 μM of working solution.
Note: Please adjust the concentration of HKOH-1 working solution according to the actual situation.
2. Cell staining
 - 2.1 Suspension cells (6-well plate)
 - a. Centrifuge at 1000 g at 4℃ for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. The cell density is 1×10⁶/mL
 - b. Add 1 mL of working solution, and then incubate at room temperature for 5-30 minutes.
 - c. Centrifuge at 400 g at 4℃ for 3-4 minutes and then discard the supernatant.

- d. Wash twice with PBS, 5 minutes each time.
 - e. Resuspend cells with serum-free cell culture medium or PBS. Observation by fluorescence microscopy or flow cytometry.
- 2.2 Adherent cells
- a. Culture adherent cells on sterile coverslips.
 - b. Remove the coverslip from the medium and aspirate excess medium.
 - c. Add 100 μ L of working solution, gently shake it to completely cover the cells, and then incubate at room temperature for 5-30 minutes.
 - d. Wash twice with medium, 5 minutes each time. Observation by fluorescence microscopy or flow cytometry.
- MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Bai X, et al. HKOH-1: A Highly Sensitive and Selective Fluorescent Probe for Detecting Endogenous Hydroxyl Radicals in Living Cells. *Angew Chem Int Ed Engl.* 2017 Oct 9;56(42):12873-12877.
- [2]. Xiaoyu Bai, et al. HKOH-1: A Highly Sensitive and Selective Fluorescent Probe for Detecting Endogenous Hydroxyl Radicals in Living Cells. *Angew Chem Int Ed Engl.* 2017 Oct 9;56(42):12873-12877.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA