


L-Ornithine

Cat. No.:	HY-B1352
CAS No.:	70-26-8
Molecular Formula:	C ₅ H ₁₂ N ₂ O ₂
Molecular Weight:	132.16
Target:	Endogenous Metabolite
Pathway:	Metabolic Enzyme/Protease
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 50 mg/mL (378.33 mM; Need ultrasonic)
DMSO : < 1 mg/mL (insoluble or slightly soluble)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	7.5666 mL	37.8329 mL	75.6659 mL
	5 mM	1.5133 mL	7.5666 mL	15.1332 mL
	10 mM	0.7567 mL	3.7833 mL	7.5666 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

L-Ornithine ((S)-2,5-Diaminopentanoic acid) is a non-proteinogenic amino acid, is mainly used in urea cycle removing excess nitrogen in vivo. L-Ornithine shows nephroprotective^{[1][2]}.

IC₅₀ & Target

Human Endogenous Metabolite

Microbial Metabolite

In Vitro

L-Ornithine (0-10 mM) activates the CaSR in a concentration-dependent manner and exerts effects on Ca²⁺ signaling in HK-2 cells^[2].

L-Ornithine (10 mM) activates TRPC channels mediated Ca²⁺ entry pathway in HK-2 cells^[2].

L-Ornithine (100 or 300 μM; 24 h) shows the protection against ROS generation in HK-2 cells via ROCE pathway^[2].

L-Ornithine (10 μM) can protect the ROS and oxidative damage in HK-2 cells exposed to H₂O₂^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Molecules. 2023 Apr 11, 28(8), 3375.
- ChemistrySelect. 2023 Jan 12.
- bioRxiv. 2023 Jun 3.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Demura S, et al. Effect of L-ornithine hydrochloride ingestion on intermittent maximal anaerobic cycle ergometer performance and fatigue recovery after exercise. Eur J Appl Physiol. 2011 Nov;111(11):2837-43.

[2]. Shin S, et al. L-ornithine activates Ca²⁺ signaling to exert its protective function on human proximal tubular cells. Cell Signal. 2020 Mar;67:109484.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA