Lamotrigine-13C3,d3

BIOLOGICAL ACTIVITY

Description

Cat. No.:	HY-B0495S	
CAS No.:	1246815-13-3	D
Molecular Formula:	$C_{6}^{13}C_{3}H_{4}D_{3}Cl_{2}N_{5}$	D
Molecular Weight:	262.09	
Target:	Sodium Channel; Autophagy	CI ¹³ Ç ^{-N} N
Pathway:	Membrane Transporter/Ion Channel; Autophagy	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	

Lamotrigine-13C3,d3 (LTG-13C3,d3) is the 13C-labeled Lamotrigine. Lamotrigine (BW430C) is a potent and orally active anticonvulsant or antiepileptic agent. Lamotrigine selectively blocks voltage-gated Na ⁺ channels, stabilizing presynaptic neuronal membranes and inhibiting glutamate release. Lamotrigine can be used for the research of epilepsy, focal seizure, et al ^{[1][2]} .

In Vitro Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

[2]. M J Leach, et al. Pharmacological studies on lamotrigine, a novel potential antiepileptic drug: II. Neurochemical studies on the mechanism of action. Epilepsia. Sep-Oct 1986;27(5):490-7.

[3]. Damianka P Getova, et al. A study of the effects of lamotrigine on mice using two convulsive tests. Folia Med (Plovdiv). Apr-Jun 2011;53(2):57-62.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898Fax: 609-228-5909E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Product Data Sheet