Marbofloxacin hydrochloride

Cat. No.:	HY-B0126A	
CAS No.:	115551-26-3	O O
Molecular Formula:	C ₁₇ H ₂₀ CIFN ₄ O ₄	FОН
Molecular Weight:	398.82	
Target:	Bacterial; Antibiotic	
Pathway:	Anti-infection	
Storage:	Please store the product under the recommended conditions in the Certificate of	HCI
	Analysis.	

BIOLOGICAL ACTIVITY		
Description	Marbofloxacin hydrochloride is a third generation fluoroquinolone and orally active antimicrobial agent, which has a broad spectrum bactericidal activity and good efficacy. Marbofloxacin hydrochloride can be used for the research of infections by Gram-positive and Gram-negative bacteria and Mycoplasma ^{[1][2][3]} .	
IC ₅₀ & Target	Quinolone	
In Vivo	Administration of Marbofloxacin at 6 mg/kg once daily for 7 days in a Staphylococcus aureus infection in tissue cages in ponies is not effective for the elimination of S. aureus infections from secluded sites [2]. The pharmacokinetic properties of marbofloxacin were investigated in 6 horses after i.v., subcutaneous and oral administration of a single dose of 2 mg/kg bwt and the minimal inhibitory concentrations (MIC) assessed for bacteria isolated from equine infectious pathologies. The clearance of marbofloxacin was mean +/- s.d. 0.25 +/- 0.05 l/kg/h and the terminal half-life 756 +/- 1.99 h. The marbofloxacin absolute bioavailabilities after subcutaneous and oral administration were 98 +/- 11% and 62 +/- 8%, respectively. Considering the breakpoint values of efficacy indices for fluoroquinolones, a marbofloxacin dosage regimen of 2 mg/kg bwt/24 h by i.v., subcutaneous or oral routes was more appropriate for enterobacteriaceae than for S. aureus [3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

REFERENCES

[1]. Shen J, et al. Marbofloxacin. Acta Crystallogr Sect E Struct Rep Online. 2012 Apr 1;68(Pt 4):0998-9.

[2]. Voermans M, et al. Clinical efficacy of intravenous administration of marbofloxacin in a Staphylococcus aureus infection in tissue cages in ponies. J Vet Pharmacol Ther. 2006 Dec;29(6):555-60.

[3]. Bousquet-Melou A, et al. Pharmacokinetics of marbofloxacin in horses. Equine Vet J. 2002 Jul;34(4):366-72.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Product Data Sheet



Page 1 of 1