


## Methimazole

<b>Cat. No.:</b>	HY-B0208		
<b>CAS No.:</b>	60-56-0		
<b>Molecular Formula:</b>	C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> S		
<b>Molecular Weight:</b>	114.17		
<b>Target:</b>	Others		
<b>Pathway:</b>	Others		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 100 mg/mL (875.89 mM)  
 H<sub>2</sub>O : ≥ 50 mg/mL (437.94 mM)  
 \* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	8.7589 mL	43.7943 mL	87.5887 mL
	5 mM	1.7518 mL	8.7589 mL	17.5177 mL
	10 mM	0.8759 mL	4.3794 mL	8.7589 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: PBS  
Solubility: 100 mg/mL (875.89 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.5 mg/mL (21.90 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (21.90 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 2.5 mg/mL (21.90 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

Methimazole is an antithyroid compound widely used for the research of hyperthyroidism. Methimazole has potent hepatotoxicity<sup>[1][2]</sup>.

## In Vitro

Methimazole inhibits CXC chemokine ligand 10 secretion in human thyrocytes<sup>[3]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Biosci Rep. 2021 May 10;BSR20204250.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

- [1]. Nakamura H, et al. Comparison of methimazole and propylthiouracil in patients with hyperthyroidism caused by Graves' disease. J Clin Endocrinol Metab. 2007 Jun;92(6):2157-62. Epub 2007 Mar 27.
- [2]. Crescioli C, et al. Methimazole inhibits CXC chemokine ligand 10 secretion in human thyrocytes. J Endocrinol. 2007 Oct;195(1):145-55.
- [3]. Tatara MR, et al. Effects of long-term oral administration of methimazole on femur and tibia properties in male Wistar rats. Biomed Pharmacother. 2017 Oct;94:124-128.
- [4]. Reza Heidari, et al. Mechanisms of methimazole cytotoxicity in isolated rat hepatocytes. Drug Chem Toxicol. 2013 Oct;36(4):403-11.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA