


## Plicamycin

<b>Cat. No.:</b>	HY-A0122
<b>CAS No.:</b>	18378-89-7
<b>Molecular Formula:</b>	C <sub>52</sub> H <sub>76</sub> O <sub>24</sub>
<b>Molecular Weight:</b>	1085.15
<b>Target:</b>	DNA/RNA Synthesis; Bacterial; Antibiotic; Glutathione S-transferase
<b>Pathway:</b>	Cell Cycle/DNA Damage; Anti-infection; Metabolic Enzyme/Protease
<b>Storage:</b>	Powder    -20°C    3 years In solvent   -80°C    6 months -20°C    1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 100 mg/mL (92.15 mM)  
 \* "≥" means soluble, but saturation unknown.

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	0.9215 mL	4.6077 mL	9.2153 mL
5 mM	0.1843 mL	0.9215 mL	1.8431 mL
10 mM	0.0922 mL	0.4608 mL	0.9215 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
 Solubility: ≥ 2.5 mg/mL (2.30 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
 Solubility: ≥ 2.5 mg/mL (2.30 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

Plicamycin (Mithramycin A) is a selective specificity protein 1 (Sp1) inhibitor. Plicamycin inhibits the growth of various cancers by decreasing Sp1 protein. Plicamycin inhibits GSTM2 promoter activity and protein expression<sup>[1][2]</sup>.

#### IC<sub>50</sub> & Target

GSTM2

GSTM2

#### In Vitro

Plicamycin (Mith) decreases Sp1 protein by inducing proteasome-dependent degradation, thereby suppressing cervical cancer growth through a DR5/caspase-8/Bid signaling pathway. Plicamycin inhibits HEp-2 and KB cell growth in a concentration-dependent manner after 48 h. Apoptotic cell death is qualitatively estimated by DAPI staining for nuclear condensation and fragmentation. Plicamycin leads to significant DNA fragmentation compared to untreated controls<sup>[1]</sup>. Plicamycin (mithramycin A) (0-400 nM, 24 h) dose-dependently inhibited GSTM2 protein expression in BFTC 905 and 5637

cells; SP1 overexpression increased GSTM2 protein expression in BFTC 905 and 5637 cells<sup>[2]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

The antitumorigenic activity of Plicamycin (0.2 mg/kg/day) is determined in a xenograft model and observed reduction in tumor volume and weight. No significant mouse body weight loss is observed in Plicamycin-treatment groups, indicating that Plicamycin-associated toxicity is minimal. Plicamycin also increases TUNEL-positive cells in tumor xenografts. No notable intergroup differences are observed among organs, indicating no marked signs of systemic toxicity at the Plicamycin dose used in this study<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Cell Assay <sup>[1]</sup>

HEp-2 cells and KB cells are cultured in DMEM 100 U/mL each of Penicillin and Streptomycin and 10% FBS for HEp-2 cells and 5% FBS for KB in a humidified atmosphere containing 5% CO<sub>2</sub> at 37°C. Equal numbers of cells are seeded and allowed to attach. At 50-60% confluence, cells are treated with DMSO or indicated concentrations of Plicamycin (50, 100, and 200 nM for HEp-2 cells; 20, 40, and 80 nM for KB cells). Cell viability is determined using CellTiter 96 Aqueous One Solution Cell Proliferation Assay Kits. In brief, cells are seeded in 96-well plates and incubated with Plicamycin. After treatment, 30 µL MTS solution is added to each well and cells are incubated for 2 h at 37°C. MTS solution is analyzed using a microplate reader at 490 nm and 690 nm<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration <sup>[1]</sup>

Mice<sup>[1]</sup>

Female nude mice are used. KB cells are suspended in sterile PBS and injected subcutaneously into the right flank of mice. Mice are randomized into two groups containing five mice each and treated with 0.2 mg/kg/day of Plicamycin (i.p.) three times per week for 29 days. Control mice receive an equal volume of vehicle. After 29 days, bodies, organs and tumors are weighed and tumor volumes determined. Tumors are measured<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Mol Cancer. 2022 Mar 18;21(1):77.
- Nat Commun. 2023 Sep 21;14(1):5891.
- Nat Commun. 2023 Feb 9;14(1):731.
- Theranostics. 2022 Jan 1;12(2):842-858.
- Dev Cell. 2024 May 17:S1534-5807(24)00296-X.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Shen CH, et al. The suppressive role of phytochemical-induced glutathione S-transferase Mu 2 in human urothelial carcinoma cells. Biomed Pharmacother. 2022 Jul;151:113102.

[2]. Choi ES, et al. Modulation of specificity protein 1 by mithramycin A as a novel therapeutic strategy for cervical cancer. Sci Rep. 2014 Nov 24;4:7162.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA