


Uridine-¹³C

Cat. No.:	HY-B1449S1		
CAS No.:	201996-62-5		
Molecular Formula:	C ₈ ¹³ CH ₁₂ N ₂ O ₆		
Molecular Weight:	245.19		
Target:	Endogenous Metabolite; Nucleoside Antimetabolite/Analog		
Pathway:	Metabolic Enzyme/Protease; Cell Cycle/DNA Damage		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

H₂O : ≥ 100 mg/mL (407.85 mM)
 H₂O : 50 mg/mL (203.92 mM; Need ultrasonic)
 DMSO : 50 mg/mL (203.92 mM; Need ultrasonic)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	4.0785 mL	20.3923 mL	40.7847 mL
	5 mM	0.8157 mL	4.0785 mL	8.1569 mL
	10 mM	0.4078 mL	2.0392 mL	4.0785 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Uridine-¹³C is the ¹³C labeled Uridine[1].

In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA