


Caffeic acid

Cat. No.:	HY-N0172		
CAS No.:	331-39-5		
Molecular Formula:	C ₉ H ₈ O ₄		
Molecular Weight:	180.16		
Target:	TRP Channel; Lipoxygenase; Endogenous Metabolite		
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (555.06 mM; Need ultrasonic)
 H₂O : < 0.1 mg/mL (ultrasonic) (insoluble)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	5.5506 mL	27.7531 mL	55.5062 mL
	5 mM	1.1101 mL	5.5506 mL	11.1012 mL
	10 mM	0.5551 mL	2.7753 mL	5.5506 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (13.88 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (13.88 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (11.55 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Caffeic acid is an inhibitor of both TRPV1 ion channel and 5-Lipoxygenase (5-LO).

IC₅₀ & Target

5-LO

In Vitro

Caffeic acid has inhibitory effects on histamine-induced responses and the inhibitory effect of Caffeic acid is gradually increased when the concentration used for pretreatment is increased from 0.1 to 1 mM, similar to typical dose-dependent

responses. Pretreatment of HEK293T-TRPV1 cells with 1 mM Caffeic acid results in significant inhibition of capsaicin-induced responses. When lower concentration of Caffeic acid is used, the inhibitory effect for capsaicin-induced responses is less evident. Calcium imaging experiments show that Caffeic acid incubation results in significant inhibition in histamine-sensitive dorsal root ganglion (DRG) neurons. Pretreatment with Caffeic acid (1 mM) results in a significant decrease in the percentage of responsive DRG neurons to histamine application from 12.5% to 2.1%. Pretreatment with 1 mM Caffeic acid dramatically blocks the allylthiocyanate (AITC)-induced intracellular calcium increase in TRPA1-expressing cells. Caffeic acid is also able to block the AITC-induced activation of TRPA1^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Mice pretreated with Caffeic acid (500 mg/kg) exhibit significantly less histamine-induced scratching (30.50±10.87 bouts/1 h, n=6). It is further found that the lower dose of Caffeic acid (100 mg/kg) is not significantly effective in terms of anti-scratching effects in histamine-induced scratching, although there appears to be a tendency of reduction (49.40±12.35 bouts/1 h, n=5). The chloroquine induced scratching is significantly inhibited by pretreatment with 500 mg/kg of Caffeic acid (161.6±31.42 bouts/1 h, n=5)^[1]. Caffeic acid significantly reduces the expression of 5-LO mRNA (P<0.01) dose-dependently in hippocampus. Compare with the ischemia-reperfusion (I/R) non-treated group, 5-LO protein expression is significantly reduced in the I/R-Caffeic acid group (P<0.05 or P<0.01), especially in the I/R-Caffeic acid group (50 mg/kg). Compare with the I/R non-treated group, the latency to find platform is significantly shortened in low- and high-dose Caffeic acid groups, the shortened platform latency is most evident in the I/R- Caffeic acid group (50 mg/kg) (P<0.01). In the low-dose Caffeic acid group, cell injury is still marked, the pyknosis ratio is (63.6±2.8)%, whereas in the high-dose Caffeic acid group, hippocampal neuron karyopyknosis is significantly reduced and the pyknosis ratio is (13.3±3.0)%^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

To determine cell viability, MTT assay is performed. HEK293T cells are cultured in 96-well plate at 37°C, a day before so that the confluence of cell is 85 to 90% on the actual day of the experiment. On the day of the experiment the cells are treated with different concentration of Caffeic acid for 10 min. Control cells are treated only with media. After removing supernatant and washing with PBS, MTT reagent (5 mg/mL) is added directly to fresh media. Cells are then incubated at 37°C for additional 4 h followed by draining of the media and overnight storage in dark condition. The next day, DMSO is added to each well and mixed in shaker for 10 min after which plate is read using microplate recorder at 490 nm with a reference wavelength of 620 nm. The relative cell viability (%) is expressed as a percentage relative to the untreated control cells^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[2]

Rats are divided into five groups: the sham group (n=9), ischemia-reperfusion (I/R) non-treated group (n=9), I/R-Caffeic acid group (10 mg/kg) (n=9), I/R-Caffeic acid group (30 mg/kg) (n=9) and I/R- Caffeic acid group (50 mg/kg) (n=9). In I/R-Caffeic acid groups, the rats are administrated Caffeic acid at 10, 30, 50 mg/kg (prepared with 0.3% sodium carboxymethyl cellulose) by intraperitoneal injection at 30 min prior to ischemia. The sham group and I/R group are treated with an equal volume of 0.3% sodium carboxymethyl cellulose^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Food Chem. 2022: 134807.
- Talanta. 2020 Feb 1;208:120450.
- Exp Cell Res. 2021 Nov 18;112934.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Pradhananga S, et al. Caffeic acid exhibits anti-pruritic effects by inhibition of multiple itch transmission pathways in mice. *Eur J Pharmacol.* 2015 Sep 5;762:313-21.
- [2]. Liang G, et al. The protective effect of caffeic acid on global cerebral ischemia-reperfusion injury in rats. *Behav Brain Funct.* 2015 Apr 18;11:18.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA