


Daidzin

Cat. No.:	HY-N0018		
CAS No.:	552-66-9		
Molecular Formula:	C ₂₁ H ₂₀ O ₉		
Molecular Weight:	416.38		
Target:	Mitochondrial Metabolism; Reverse Transcriptase; Aldehyde Dehydrogenase (ALDH)		
Pathway:	Metabolic Enzyme/Protease; Anti-infection		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 250 mg/mL (600.41 mM; ultrasonic and warming and heat to 60°C)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	2.4017 mL	12.0083 mL	24.0165 mL
	5 mM	0.4803 mL	2.4017 mL	4.8033 mL
	10 mM	0.2402 mL	1.2008 mL	2.4017 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 0.5% CMC-Na/1% Tween-80 in Saline water
Solubility: 39.6 mg/mL (95.11 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (5.00 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: 2.08 mg/mL (5.00 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.08 mg/mL (5.00 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Daidzin is an isoflavone with antioxidant, anticancer, and antiatherosclerotic activities. Daidzin is a potent and selective inhibitor of mitochondrial ALDH-2. Daidzin reduces ethanol consumption^[1].

In Vitro

Daidzin inhibits hamster and rat ALDH-2, with K_is of 0.082 and 0.052 μM respectively^[1]. Daidzin (500 μM) shows antioxidant activity, measured by its ability to inhibit cytochrome c reduction by Phorbol myristic

	<p>acetate (PMA)-stimulated neutrophils^[3]. Daidzin (0-30 μM, 30 min) inhibits the formation of 5-HIAA (IC₅₀ value of \approx2.7 μM), and increases the amounts of 5-HIAL and 5-HTOL in the incubation media of hamster liver mitochondrial^[4]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
In Vivo	<p>Daidzin (30 mg/kg, intragastrically) decreases blood alcohol levels (induced by 3 g ethanol/kg body weight), and shortens sleep time induced by Ethanol intoxication (7 g/kg by intubation) in fasted rats^[3]. Daidzin (70 meq/hamster/day, i.p.) inhibits hamster ethanol intake by 62%^[4]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

CUSTOMER VALIDATION

- Free Radical Bio Med. 2020 May 20;152:8-17.
- J Ethnopharmacol. 2022 Aug 13;115593.
- Mol Med Rep. 2020 Sep;22(3):2373-2385.
- Biochem Biophys Res Commun. 2023 Mar 24;657:108-118.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Keung WM, Vallee BL. Daidzin and its antidipsotropic analogs inhibit serotonin and dopamine metabolism in isolated mitochondria. Proc Natl Acad Sci U S A. 1998 Mar 3;95(5):2198-203.
- [2]. Xie Cl, et al. Daidzin, an antioxidant isoflavonoid, decreases blood alcohol levels and shortens sleep time induced by ethanol intoxication. Alcohol Clin Exp Res. 1994 Dec;18(6):1443-7.
- [3]. Keung WM, et al. Kudzu root: an ancient Chinese source of modern antidipsotropic agents. Phytochemistry. 1998 Feb;47(4):499-506.
- [4]. Keung WM, et al. Daidzin inhibits mitochondrial aldehyde dehydrogenase and suppresses ethanol intake of Syrian golden hamsters. Proc Natl Acad Sci U S A. 1997 Mar 4;94(5):1675-9.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA