**Proteins** 

## (E/Z)-Polydatin

Cat. No.: HY-N0120 CAS No.: 65914-17-2 Molecular Formula:  $C_{20}H_{22}O_8$ Molecular Weight: 390.38 Others Target: Pathway: Others

Storage: Powder 3 years 2 years

-80°C In solvent 2 years

-20°C

-20°C 1 year

**Product** Data Sheet

## **SOLVENT & SOLUBILITY**

In Vitro

DMSO: 150 mg/mL (384.24 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	2.5616 mL	12.8080 mL	25.6161 mL
	5 mM	0.5123 mL	2.5616 mL	5.1232 mL
	10 mM	0.2562 mL	1.2808 mL	2.5616 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.40 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.40 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.40 mM); Clear solution

## **BIOLOGICAL ACTIVITY**

Description

(E/Z)-Polydatin ((E/Z)-Piceid) is a monocrystalline compound originally isolated from the root and rhizome of Polygonum cuspidatum. (E/Z)-Polydatin has anti-platelet aggregation, anti-oxidative action of low-density lipoprotein (LDL), cardioprotective activity, anti-inflammatory and immune-regulating functions<sup>[1]</sup>.

## **REFERENCES**

[1]. Du QH, et al. Polydatin: a review of pharmacology and pharmacokinetics. Pharm Biol. 2013 Nov;51(11):1347-54.							
Caution: Product has not been fully validated for medical applications. For research use only.							
	Tel: 609-228-6898	Fax: 609-228-5909  Deer Park Dr. Suite O. Monm	E-mail: tech@MedChemExpress.outh Junction, NJ 08852, USA	com			
	71441.0001.2	Joen am J., Jane Q. monni					

Page 2 of 2 www.MedChemExpress.com