


Ginkgolide B

Cat. No.:	HY-N0784		
CAS No.:	15291-77-7		
Molecular Formula:	C ₂₀ H ₂₄ O ₁₀		
Molecular Weight:	424.4		
Target:	Apoptosis; Platelet-activating Factor Receptor (PAFR)		
Pathway:	Apoptosis; GPCR/G Protein		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 100 mg/mL (235.63 mM)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		2.3563 mL	11.7813 mL	23.5627 mL
	5 mM		0.4713 mL	2.3563 mL	4.7125 mL
	10 mM		0.2356 mL	1.1781 mL	2.3563 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 0.5% CMC-Na/saline water
Solubility: 5 mg/mL (11.78 mM); Suspended solution; Need ultrasonic and warming and heat to 44°C
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (5.89 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (4.90 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.08 mg/mL (4.90 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Ginkgolide B (BN-52021), a terpene lactone, is a potent platelet activating factor antagonist. Ginkgolide B protects endothelial cells via the activation of PXR from injuries induced by xeno- and endobiotics. Ginkgolide B can pass through the brain-blood barrier. Ginkgolide B has anti-oxidant, anti-inflammatory, anti-tumor, and anti-apoptotic activity. Ginkgolide B has marked neuroprotective effects against ischemia-induced impairments^{[1][2]}.

In Vitro	<p>Ginkgolide B (BN-52021; 30-150 μM; 24 h) inhibits the up-regulation of VCAM-1 and E-selectin protein expression induced by TNF-α (10 ng/mL)^[1].</p> <p>Ginkgolide B (30-150 μM; 24, 48 h) increases CYP3A4 mRNA and MDR1 expression^[1].</p> <p>Ginkgolide B (30-150 μM; pretreated for 24 h) reduces Staurosporine-induced (500 nM; 6 h) apoptosis compared with the positive controls, which exhibited marked apoptosis compared with negative controls. Ginkgolide B does not affect Doxorubicin (5 μM; 24 h) induced apoptosis^[1].</p> <p>Ginkgolide B regulates PXR activity and does not affect PXR expression^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Western Blot Analysis^[1]</p>	
	Cell Line:	HUVECs
	Concentration:	30, 60, 150 μ M
	Incubation Time:	24 h
	Result:	Inhibited the up-regulation of VCAM-1 and E-selectin protein expression induced by TNF- α .
	RT-PCR ^[1]	
	Cell Line:	HUVECs
	Concentration:	30, 60, 150 μ M
	Incubation Time:	24, 48 h
	Result:	Increased CYP3A4 mRNA and MDR1 expression after 24 h, and this effect was dose-dependent after 48 h.
In Vivo	<p>Ginkgolide B (BN-52021; 20 mg/kg; Intraperitoneally; immediately and 6 hours after ischemia, and thereafter once daily; 14 days) decreases the neurological deficit score^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>	
	Animal Model:	Male clean healthy Sprague-Dawley rats, weighing 250-280 g and aged 8 weeks ^[2]
	Dosage:	20 mg/kg
	Administration:	Intraperitoneally; immediately and 6 hours after ischemia, and thereafter once daily; 14 days
	Result:	Decreased the neurological deficit score, increased the proportion of nestin-, neuron-specific enolase- and glial fibrillary acid protein-positive cells. Caused down-regulation of NSE protein at day 14.

CUSTOMER VALIDATION

- Biofactors. 2019 Nov;45(6):950-958.
- Neuropharmacology. 2024 Mar 14;250:109907.
- Research Square Print. 2022 Jun.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Tao Zhou, et al. Ginkgolide B protects human umbilical vein endothelial cells against xenobiotic injuries via PXR activation. *Acta Pharmacol Sin.* 2016 Feb;37(2):177-86.
- [2]. Pei-Dong Zheng, et al. Ginkgolide B promotes the proliferation and differentiation of neural stem cells following cerebral ischemia/reperfusion injury, both in vivo and in vitro. *Neural Regen Res.* 2018 Jul;13(7):1204-1211.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA