


## L-Glutamine-5-<sup>13</sup>C

<b>Cat. No.:</b>	HY-N0390S4		
<b>CAS No.:</b>	159680-32-7		
<b>Molecular Formula:</b>	C <sub>4</sub> <sup>13</sup> CH <sub>10</sub> N <sub>2</sub> O <sub>3</sub>		
<b>Molecular Weight:</b>	147.14		
<b>Target:</b>	mGluR; Ferroptosis; Endogenous Metabolite		
<b>Pathway:</b>	GPCR/G Protein; Neuronal Signaling; Apoptosis; Metabolic Enzyme/Protease		
<b>Storage:</b>	Powder	-20°C	3 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

H<sub>2</sub>O : 25 mg/mL (169.91 mM; ultrasonic and warming and heat to 60°C)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	6.7962 mL	33.9812 mL	67.9625 mL
5 mM	1.3592 mL	6.7962 mL	13.5925 mL
10 mM	0.6796 mL	3.3981 mL	6.7962 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

L-Glutamine-5-<sup>13</sup>C is the <sup>13</sup>C-labeled L-Glutamine. L-Glutamine (L-Glutamic acid 5-amide) is a non-essential amino acid present abundantly throughout the body and involved in many metabolic processes. L-Glutamine provides a source of carbons for oxidation in some cells[1][2].

#### In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA