L-Serine-¹³C₃, ¹⁵N

Cat. No.: HY-N0650S5 202407-34-9 CAS No.: ¹³C₃H₇ ¹⁵NO₃ Molecular Formula:

Molecular Weight: 109.06

Target: **Endogenous Metabolite** Pathway: Metabolic Enzyme/Protease

Storage: 4°C, sealed storage, away from moisture and light

* In solvent: -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture

and light)

$$H_{2}$$
 H_{2}
 H_{3}
 H_{3}
 H_{3}
 H_{3}
 H_{4}
 H_{5}
 H_{5}
 H_{5}
 H_{5}
 H_{6}
 H_{7}
 H_{7

Product Data Sheet

SOLVENT & SOLUBILITY

In Vitro

H₂O: 50 mg/mL (458.46 mM; Need ultrasonic) H₂O: 50 mg/mL (458.46 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	9.1693 mL	45.8463 mL	91.6926 mL
	5 mM	1.8339 mL	9.1693 mL	18.3385 mL
	10 mM	0.9169 mL	4.5846 mL	9.1693 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

L-Serine-¹³C₃, ¹⁵N is the ¹³C- and ¹⁵N-labeled L-Serine. L-Serine ((-)-Serine; (S)-Serine), one of the so-called non-essential Description amino acids, plays a central role in cellular proliferation.

In Vitro Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to

affect the pharmacokinetic and metabolic profiles of drugs^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com