


Tectorigenin

Cat. No.:	HY-N0792		
CAS No.:	548-77-6		
Molecular Formula:	C ₁₆ H ₁₂ O ₆		
Molecular Weight:	300.26		
Target:	Others		
Pathway:	Others		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 120 mg/mL (399.65 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM	3.3304 mL	16.6522 mL	33.3045 mL
		5 mM	0.6661 mL	3.3304 mL	6.6609 mL
10 mM		0.3330 mL	1.6652 mL	3.3304 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	<ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (6.93 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (6.93 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (6.93 mM); Clear solution 				

BIOLOGICAL ACTIVITY

Description	Tectorigenin is a plant isoflavonoid originally isolated from the dried flower of <i>Pueraria lobate</i> Benth.
In Vitro	Tectorigenin is a plant isoflavonoid originally isolated from the dried flower of <i>Pueraria thomsonii</i> Benth. Palmitic acid (PA)-stimulated ROS production is abolished by treatment with Tectorigenin for HUVECs in a dose-dependent manner (0.1, 1, 10 μM). Treatment with Tectorigenin attenuates enhanced IKKβ phosphorylation and effectively blocks NF-κB activation by inhibition of p65 phosphorylation at concentrations ranging from 0.1 to 10 μM. Tectorigenin treatment also effectively inhibits PA-augmented TNF-α and IL-6 production in a concentration dependent manner ^[1] . The number of viable HepG2 cells treated by Tectorigenin decreases in a concentration- and time-dependent manner. When HepG2 cells are treated with

Tectorigenin at 5, 10 and 20 mg/L for 24 h, the viability rate is 91%, 79% and 62%, respectively^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

HUVECs grown to confluence in 24-well plates are pretreated with Tectorigenin (0.1, 1, 10 μ M), salicylate (5 mM) or GSH (1 mM) for 30 min, then stimulated with Palmitic acid (PA) (100 μ M) for further 12 h in serum-free medium, and the medium is then collected on ice. The levels of TNF- α and IL-6 in the supernatant are assayed with commercial ELISA Kits^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Assay ^[2]

Cell viability is assessed by MTT method. Briefly, cells are seeded in 96-well plate at a density of 1×10^4 cells/well. After 24 h incubation, Tectorigenin at different concentrations is added to the cells while only DMSO (solvent) is added as a negative control. After growing for 12, 24 and 48 h, cells are incubated with MTT (0.5 mg/mL) for 4 h at 37°C. During this incubation period, water-insoluble formazan crystals are formed, which are dissolved by the addition of 100 μ L/well DMSO. The optical densities at 570 nm are measured using an enzyme-linked immunosorbent assay plate reader. Wells containing culture medium and MTT but no cells act as blanks^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Br J Pharmacol. 2021 Jun;178(12):2443-2460.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Wang Q, et al. Tectorigenin Attenuates Palmitate-Induced Endothelial Insulin Resistance via Targeting ROS-Associated Inflammation and IRS-1 Pathway. PLoS One. 2013 Jun 19;8(6):e66417.

[2]. Jiang CP, et al. Pro-apoptotic effects of tectorigenin on human hepatocellular carcinoma HepG2 cells. World J Gastroenterol. 2012 Apr 21;18(15):1753-64.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA