


Vitamin K1

Cat. No.:	HY-N0684		
CAS No.:	84-80-0		
Molecular Formula:	C ₃₁ H ₄₆ O ₂		
Molecular Weight:	450.7		
Target:	Endogenous Metabolite		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

Ethanol : ≥ 50 mg/mL (110.94 mM)
 DMSO : 5.6 mg/mL (12.43 mM; Need ultrasonic and warming)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		2.2188 mL	11.0939 mL	22.1877 mL
	5 mM		0.4438 mL	2.2188 mL	4.4375 mL
	10 mM		0.2219 mL	1.1094 mL	2.2188 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% EtOH >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 2.5 mg/mL (5.55 mM); Clear solution
- Add each solvent one by one: 10% EtOH >> 90% (20% SBE-β-CD in saline)
 Solubility: 2.5 mg/mL (5.55 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% EtOH >> 90% corn oil
 Solubility: ≥ 2.5 mg/mL (5.55 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Vitamin K1 a naturally occurring vitamin required for blood coagulation and bone and vascular metabolism.

IC₅₀ & Target

Human Endogenous Metabolite

In Vitro

Phylloquinone (Vitamin K1) is a prenylated naphthoquinone that is synthesized exclusively by plants, green algae, and some

species of cyanobacteria, where it serves as a vital electron carrier in photosystem I and as an electron acceptor for the formation of protein disulfide bonds. In humans and other vertebrates, phyloquinone plays the role of a vitamin (vitamin K1) that is required for blood coagulation and bone and vascular metabolism. Phyloquinone from green leafy vegetables and vegetable oil represents the major dietary source of vitamin K for humans^[1]. Vitamin K1 treatment causes a significant antiproliferative effect and induces apoptosis in Caco-2, HT-29, and SW480 cell lines, with the involvement of the MAPK pathway. A concomitant and significant decrease in the polyamine biosynthesis occurs^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Subjects who increase their dietary intake of vitamin K during the follow-up had a 51% reduced risk of incident diabetes compared with subjects who decrease or does not change the amount of phyloquinone intake^[3]. Vitamin K supplementation reverses the high fat diet induced bone deterioration by modulating osteoblast and osteoclast activities and prevent bone loss in a high-fat diet-induced obese mice^[4]. Application of vitamin K1 to the skin has been used for suppression of pigmentation and resolution of bruising. The effects produced by the topical vitamin K1 shows significant healing when compared with control group in parameters such as wound contraction, epithelialization period, hydroxyproline content and tensile strength^[5].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[2]

Caco-2, HT-29, and SW480 cells are treated with increasing concentrations of vitamin K1 (10, 50, 100, and 200 μ M) for 24 h, 48 h, and 72 h. MTT is added to each dish and incubated for 2 h at 37°C. At the end of the incubation period, the medium is removed. The plate is read at 570 nm^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^{[4][5]}

Rats: For inducing full-thickness wound in rats, the excisional wound model is used. Five groups consisting of 8 rats each are used. Vitamin K cream (1% and 2%, w/w) is prepared in eucerin base and applied on the wound once a day until complete healing had occurred. Healing is defined by decreased wound margin (wound contraction), re-epithelialization, tensile strength and hydroxyproline content. Histopathological examination is also done^[5].

Mice: Four-week-old C57BL/6J male mice are fed a 10% fat normal diet group or a 45% kcal high-fat diet group, with or without 200 mg/1000 g vitamin K1 (Normal diet + K1, high-fat diet + K1) and 200 mg/1000 g vitamin K2 (Normal diet + K2, high-fat diet + K2) for 12 weeks^[4].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Med Chem. 2021 Sep 21.
- Int J Mol Sci. 2023 Apr 28, 24(9), 8012.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Basset GJ, et al. Phyloquinone (Vitamin K1): Occurrence, Biosynthesis and Functions. Mini Rev Med Chem. 2016 Jun 22.

[2]. Orlando A, et al. Vitamin K1 exerts antiproliferative effects and induces apoptosis in three differently graded human colon cancer cell lines. Biomed Res Int. 2015;2015:296721.

[3]. Ibarrola-Jurado N, et al. Dietary phyloquinone intake and risk of type 2 diabetes in elderly subjects at high risk of cardiovascular disease. Am J Clin Nutr. 2012

Nov;96(5):1113-8.

[4]. Kim M, et al. Vitamin K1 (phylloquinone) and K2 (menaquinone-4) supplementation improves bone formation in a high-fat diet-induced obese mice. J Clin Biochem Nutr. 2013 Sep;53(2):108-13.

[5]. Hemmati AA, et al. Topical vitamin K1 promotes repair of full thickness wound in rat. Indian J Pharmacol. 2014 Jul-Aug;46(4):409-12.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA